arXiv:0804.1153v2 [quant-ph] 5 Jun 2008

Groups of Operators for Evolution Equations of Quantum Many-Particle Systems

V.I. Gerasimenko*

Institute of Mathematics of NAS of Ukraine, Tereshchenkivs'ka Str., 3,
01601 Kyiv-4, Ukraine
gerasym@imath.kiev.ua

Abstract

The aim of this work is to study the properties of groups of operators for evolution equations of quantum many-particle systems, namely, the von Neumann hierarchy for correlation operators, the BBGKY hierarchy for marginal density operators and the dual BBGKY hierarchy for marginal observables. We show that the concept of cumulants (semi-invariants) of groups of operators for the von Neumann equations forms the basis of the expansions for one-parametric families of operators for evolution equations of infinitely many particles.

MSC(2000): 35Q40; 47d06.
Keywords. Quantum dynamical semigroup, quantum many-particle system, cluster expansion, cumulant (semi-invariant), BBGKY hierarchy, dual BBGKY hierarchy, von Neumann hierarchy.

1 Introduction

Recently we observe significant progress in the study of the evolution equations of quantum many-particle systems [1],[2]. In particular it is involved in such fundamental problem as the rigorous derivation of quantum kinetic equations [3]-[9].

As is well known, there are various possibilities to describe the evolution of quantum many-particle systems [1]. The sequence of the von Neumann equations for density operators [2],[11], the von Neumann hierarchy for correlation operators [23], the BBGKY hierarchy for marginal density operators [11] and the dual BBGKY hierarchy for marginal observables [1] give the equivalent approaches for the description of the evolution of finitely many particles. Papers [23]-[27] constructed the one-parametric families of operators that define solutions of the Cauchy problem for these evolution equations. It was established that the concept of cumulants (semi-invariants) of groups of operators for the von Neumann equations forms the basis of the one-parametric families of operators of various evolution equations of quantum systems of particles, in particular, the BBGKY hierarchy for infinitely many particles [23].

The aim of the paper is to investigate properties of groups of operators for evolution equations of quantum many-particle systems related with their cumulant structure on suitable Banach spaces.

In the beginning we will formulate some necessary facts about the description of quantum many-particle systems.

---

*Partially supported by the WTZ grant No M/124 (UA 04/2007)

1.1 Group of Operators for the von Neumann Equation

Let a sequence $f = \big(I, f_1, \ldots, f_n, \ldots\big)$ is an infinite sequence of self-adjoint operators f_n (I is a unit operator) defined on the Fock space $\mathcal{F}_{\mathcal{H}} = \bigoplus_{n=0}^{\infty}\mathcal{H}^{\otimes n}$ over the Hilbert space $\mathcal{H}$ ($\mathcal{H}^0 = \mathbb{C}$). Operators f_n defined in the n-particle Hilbert space $\mathcal{H}_n = \mathcal{H}^{\otimes n}$ we will denote by $f_n(1, \ldots, n)$. For a system of identical particles obeying Maxwell-Boltzmann statistics, one has $f_n(1, \ldots, n) = f_n(i_1, \ldots, i_n)$ if $\{i_1, \ldots, i_n\} \in \{1, \ldots, n\}$.

Let $\mathfrak{L}^1_\alpha(\mathcal{F}_{\mathcal{H}}) = \bigoplus_{n=0}^{\infty}\alpha^n \mathfrak{L}^1(\mathcal{H}_n)$ be the space of sequences $f = \big(I, f_1, \ldots, f_n, \ldots\big)$ of trace class operators $f_n = f_n(1, \ldots, n) \in \mathfrak{L}^1(\mathcal{H}_n)$, satisfying the above-mentioned symmetry condition, equipped with the trace norm

$$\|f\|_{\mathfrak{L}^1_\alpha(\mathcal{F}_{\mathcal{H}})} = \sum_{n=0}^{\infty} \alpha^n \|f_n\|_{\mathfrak{L}^1(\mathcal{H}_n)} = \sum_{n=0}^{\infty} \alpha^n \ \mathrm{Tr}_{1,\ldots,\mathrm{n}} |f_n(1, \ldots, n)|,$$

where $\alpha > 1$ is a real number, $\mathrm{Tr}_{1,\ldots,\mathrm{n}}$ is the partial trace over $1, \ldots, n$ particles. We will denote by $\mathfrak{L}^1_{\alpha,0}$ the everywhere dense set in $\mathfrak{L}^1_\alpha(\mathcal{F}_{\mathcal{H}})$ of finite sequences of degenerate operators [14] with infinitely differentiable kernels with compact supports. We will also consider the space $\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}}) = \bigoplus_{n=0}^{\infty}\mathfrak{L}^1(\mathcal{H}_n)$.

We note that the sequences of operators $f_n \in \mathfrak{L}^1(\mathcal{H}_n)$, $n \geq 1$, whose kernels are known as density matrices [12] defined on the n-particle Hilbert space $\mathcal{H}_n = \mathcal{H}^{\otimes n} = L^2(\mathbb{R}^{\nu n})$, describe the states of a quantum system of non-fixed number of particles. The space $\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$ contains sequences of operators more general than those determining the states of systems.

The evolution of all possible states of quantum systems is described by the initial-value problem to the von Neumann equation [10, 11]. A solution of such Cauchy problem is defined by the following one-parametric family of operators on $\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$

$$\mathbb{R}^1 \ni t \mapsto \mathcal{G}(-t)f := \mathcal{U}(-t)f\mathcal{U}^{-1}(-t), \tag{1.1}$$

where $f \in \mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$ and $\mathcal{U}(-t) = \bigoplus_{n=0}^{\infty}\mathcal{U}_n(-t)$,

$$\begin{aligned} \mathcal{U}_n(-t) &:= e^{-\frac{i}{\hbar}tH_n}, \\ \mathcal{U}_n^{-1}(-t) &:= e^{\frac{i}{\hbar}tH_n}, \end{aligned} \tag{1.2}$$

$\mathcal{U}_0(-t) = I$ is a unit operator. The Hamiltonian $H = \bigoplus_{n=0}^{\infty}H_n$ in (1.2) is a self-adjoint operator with domain $\mathcal{D}(H) = \{\psi = \oplus_{n=0}^{\infty}\psi_n \in \mathcal{F}_{\mathcal{H}} \mid \psi_n \in \mathcal{D}(H_n) \subset \mathcal{H}_n, \ \sum_n \|H_n\psi_n\|^2 < \infty\} \subset \mathcal{F}_{\mathcal{H}}$ [14].

Assume $\mathcal{H} = L^2(\mathbb{R}^3)$ then an element $\psi \in \mathcal{F}_{\mathcal{H}} = \bigoplus_{n=0}^{\infty}L^2(\mathbb{R}^{3n})$ is a sequence of functions $\psi = \big(\psi_0, \psi_1(q_1), \ldots, \psi_n(q_1, \ldots, q_n), \ldots\big)$ such that $\|\psi\|^2 = |\psi_0|^2 + \sum_{n=1}^{\infty}\int dq_1 \ldots dq_n \ |\psi_n(q_1, \ldots, q_n)|^2 < +\infty$. On the subspace of infinitely differentiable functions with compact supports $\psi_n \in L^2_0(\mathbb{R}^{3n}) \subset L^2(\mathbb{R}^{3n})$, n-particle Hamiltonian H_n acts according to the formula ($H_0 = 0$)

$$H_n\psi_n = -\frac{\hbar^2}{2}\sum_{i=1}^{n}\Delta_{q_i}\psi_n + \sum_{k=1}^{n}\sum_{i_1<\ldots<i_k=1}^{n}\Phi^{(k)}(q_{i_1}, \ldots, q_{i_k})\psi_n. \tag{1.3}$$

where $\Phi^{(k)}$ is a k-body interaction potential satisfying Kato conditions [14] and $h = 2\pi\hbar$ is a Planck constant.

We remark that the nature of notations (1.2) used for unitary groups $e^{\pm\frac{i}{\hbar}tH_n}$ is related to the correspondence principle between quantum and classical systems [1] and is a consequence of the existence of two approaches to the description of the evolution of systems in the framework of observables or states.

The properties of a one-parametric family $\{\mathcal{G}(-t)\}_{t\in\mathbb{R}}$ of operators (1.1) follow from the properties of groups (1.2) described, for example, in [12].

Proposition 1.1 ([10],[11]). *On the space $\mathfrak{L}^1(\mathcal{F}_\mathcal{H})$ mapping* (1.1) *defines an isometric strongly continuous group, i.e. one is a C_0-group, which preserves positivity and self-adjointness of operators.*

If $f \in \mathfrak{L}^1_0(\mathcal{F}_\mathcal{H}) \subset \mathcal{D}(-\mathcal{N}) \subset \mathfrak{L}^1(\mathcal{F}_\mathcal{H})$ then in the sense of the norm convergence of the space $\mathfrak{L}^1(\mathcal{F}_\mathcal{H})$ there exists a limit that is determined the infinitesimal generator: $-\mathcal{N} = \bigoplus_{n=0}^{\infty}(-\mathcal{N}_n)$ of group (1.1)

$$\lim_{t\rightarrow 0}\frac{1}{t}\big(\mathcal{G}(-t)f-f\big) = -\frac{i}{\hbar}(Hf-fH) := -\mathcal{N}f, \tag{1.4}$$

where $H = \bigoplus_{n=0}^{\infty} H_n$ is the Hamiltonian (1.3) *and the operator: $(-i/\hbar)(Hf-fH)$ is defined on the domain $\mathcal{D}(H) \subset \mathcal{F}_\mathcal{H}$.*

Group of operators (1.1) and infinitesimal generator (1.4) studied within the framework of kernels and symbols of the operators in [12] and for the Wigner representation in [19]. Some applications of evolution groups of quantum systems are considered in [18].

1.2 Group of Operators for the Heisenberg Equation

The adjoint to $\mathfrak{L}^1(\mathcal{F}_\mathcal{H})$ space is isometric to the space $\mathfrak{L}(\mathcal{F}_\mathcal{H})$ of sequences $g = \big(I, g_1, \ldots, g_n, \ldots\big)$ of bounded operators g_n (I is a unit operator) defined on the Hilbert space $\mathcal{H}_n$ satisfying symmetry property $g_n(1,\ldots,n) = g_n(i_1,\ldots,i_n)$ for $\{i_1,\ldots,i_n\} \in \{1,\ldots,n\}$ with an operator norm. The space $\mathfrak{L}(\mathcal{F}_\mathcal{H})$ is dual to the space $\mathfrak{L}^1(\mathcal{F}_\mathcal{H})$ with respect to the bilinear form

$$\big\langle g\big|f\big\rangle = \sum_{n=0}^{\infty}\frac{1}{n!}\mathrm{Tr}_{1,\ldots,\mathrm{n}}\, g_n f_n. \tag{1.5}$$

We will also consider more general space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ than $\mathfrak{L}(\mathcal{F}_\mathcal{H})$ with a norm

$$\|g\|_{\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})} = \max_{n\geq 0}\ \frac{\gamma^n}{n!}\ \|g_n\|_{\mathfrak{L}(\mathcal{H}_n)},$$

where $0<\gamma<1$ and $\|.\|_{\mathfrak{L}(\mathcal{H}_n)}$ is an operator norm [14].

An observable of finitely many quantum particles is a sequence of self-adjoint operators from $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ and positive normalized continuous linear functional (1.5) on the space of observables is interpreted as its mean value [13]. The case of the unbounded observables can be reduced to the case under consideration [10].

The evolution of observables is described by the initial-value problem to the Heisenberg equation (the dual von Neumann equation) [10, 11]. On $\mathfrak{L}(\mathcal{F}_\mathcal{H})$ a solution of the Cauchy problem to the Heisenberg equation is defined the following one-parametric family of operators [10]

$$\mathbb{R}^1 \ni t \mapsto \mathcal{G}(t)g := \mathcal{U}(t)g\mathcal{U}^{-1}(t), \tag{1.6}$$

where $\mathcal{U}(t) = \bigoplus_{n=0}^{\infty}\mathcal{U}_n(t)$ and the operators $\mathcal{U}_n(t)$, $\mathcal{U}_n^{-1}(t)$ are defined by formulas (1.2). Mapping $\mathcal{G}(t)$ (1.6) is adjoint (dual) to $\mathcal{G}(-t)$ (1.1).

Proposition 1.2 ([10],[13]). *On the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ mapping* (1.6) *defines an isometric $*$-weak continuous group, i.e. one is a C_0^*-group. This group preserves the self-adjointness of operators. The infinitesimal generator $\mathcal{N} = \bigoplus_{n=0}^{\infty}\mathcal{N}_n$ of this group of operators is a closed operator for the $*$-weak topology and on its domain of the definition $\mathcal{D}(\mathcal{N}) \subset \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ which is the everywhere dense set for the $*$-weak topology it is defined in the sense of the $*$-weak convergence of the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ by the formula*

$$\mathrm{w}^*\!-\lim_{t\rightarrow 0}\frac{1}{t}\big(\mathcal{G}(t)g-g\big) = -\frac{i}{\hbar}(gH-Hg), \tag{1.7}$$

where $H = \bigoplus_{n=0}^{\infty} H_n$ is the Hamiltonian (1.3) *and the operator: $\mathcal{N}g = (-i/\hbar)(gH-Hg)$ is defined on the domain $\mathcal{D}(H) \subset \mathcal{F}_\mathcal{H}$.*

1.3 Cumulants of Groups of Operators

Further to the group $\{\mathcal{G}(-t)\}_{t\in\mathbb{R}}$ we will also consider more general mappings on the space $\mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$. Namely let us expand the group $\mathcal{G}(-t)=\bigoplus_{n=0}^\infty \mathcal{G}_n(-t)$ as following cluster expansions

$$\mathcal{G}_n(-t,Y)=\sum_{\mathrm{P}:Y=\bigcup_i X_i}\ \prod_{X_i\subset \mathrm{P}}\mathfrak{A}_{|X_i|}(t,X_i),\qquad n=|Y|\geq 0, \tag{1.8}$$

where $\sum_\mathrm{P}$ is the sum over all possible partitions P of the set $Y\equiv(1,\ldots,n)$ into $|\mathrm{P}|$ nonempty mutually disjoint subsets $X_i\subset Y$. A solution of recurrence relations (1.8) is determined by the expansions [26]

$$\mathfrak{A}_n(t,Y)=\sum_{\mathrm{P}:Y=\bigcup_i X_i}(-1)^{|\mathrm{P}|-1}(|\mathrm{P}|-1)!\prod_{X_i\subset \mathrm{P}}\mathcal{G}_{|X_i|}(-t,X_i),\qquad n=|Y|\geq 0, \tag{1.9}$$

where the notations are similar to that in formula (1.8). The operator $\mathfrak{A}_n(t)$ we refer to as the *nth*-order cumulant (semi-invariant) of evolution operators (1.1). Some properties of cumulants (1.9) considered in [23].

The generator of the 1*st*-order cumulant is given by operator (1.4), i.e.

$$\lim_{t\rightarrow 0}\frac{1}{t}\big(\mathfrak{A}_1(t,Y)-I\big)f_n(Y)=-\mathcal{N}_n(Y)f_n(Y),$$

where for $f_n\in\mathfrak{L}^1_0(\mathcal{H}_n)$ this limit exists in the sense of the norm convergence of the space $\mathfrak{L}^1(\mathcal{H}_n)$.

The infinitesimal generator of the *nth*-order cumulant, $n\geq 2$, is an operator $(-\mathcal{N}_{\mathrm{int}}^{(n)})$ defining by n-body interaction potential (1.3). According to the equality

$$\sum_{\mathrm{P}:\,Y=\bigcup_i X_i}(-1)^{|\mathrm{P}|-1}(|\mathrm{P}|-1)!=\sum_{k=1}^{n}(-1)^{k-1}\mathrm{s}(n,k)(k-1)!=\delta_{n,1}, \tag{1.10}$$

where $\mathrm{s}(n,k)$ is the Stirling numbers of the second kind and $\delta_{n,1}$ is a Kroneker symbol, for the *nth*-order cumulant, $n\geq 2$, in the sense of a point-by-point convergence of the space $\mathfrak{L}^1(\mathcal{H}_n)$ we have

$$\lim_{t\rightarrow 0}\frac{1}{t}\,\mathfrak{A}_n(t,Y)f_n(Y)=\sum_{\mathrm{P}:\,Y=\bigcup_i X_i}(-1)^{|\mathrm{P}|-1}(|\mathrm{P}|-1)!\sum_{X_i\subset\mathrm{P}}(-\mathcal{N}_{|X_i|}(X_i))f_n(Y)=$$

$$\sum_{\mathrm{P}:\,Y=\bigcup_i X_i}(-1)^{|\mathrm{P}|-1}(|\mathrm{P}|-1)!\sum_{X_i\subset\mathrm{P}}\ \sum_{k=2}^{|X_i|}\ \sum_{i_1<\ldots<i_k\in\{X_i\}}\big(-\mathcal{N}_{\mathrm{int}}^{(k)}(i_1,\ldots,i_k)\big)f_n(Y).$$

Here for the operator $\Phi^{(n)}$ from Hamiltonian (1.3) the operator $\mathcal{N}_{\mathrm{int}}^{(n)}$ is defined by the formula

$$\mathcal{N}_{\mathrm{int}}^{(n)}f_n:=-\frac{i}{\hbar}\big(f_n\Phi^{(n)}-\Phi^{(n)}f_n\big). \tag{1.11}$$

Summing coefficients before every operator $\mathcal{N}_{\mathrm{int}}^{(k)}$ we deduce

$$\lim_{t\rightarrow 0}\frac{1}{t}\mathfrak{A}_n(t,Y)f_n(Y)=-\mathcal{N}_{\mathrm{int}}^{(n)}(Y)f_n(Y), \tag{1.12}$$

Thus for $f_n\in\mathfrak{L}^1_0(\mathcal{H}_n)$ the generator of the *nth*-order cumulant is defined by formula (1.12) in the sense of the norm convergence of the space $\mathfrak{L}^1(\mathcal{H}_n)$.

The dual cumulants of the groups $\mathcal{G}(t)=\bigoplus_{n=0}^\infty\mathcal{G}_n(t)$ (1.6) will be introduced in Section 4. For classical many-particle systems cumulants of evolution operators were introduced in [26],[27].

2 Group of Operators for the von Neumann Hierarchy

The evolution of correlations of quantum finitely many particles is described by the von Neumann hierarchy for the correlation operators [23]. It is an equivalent approach for the description of the evolution of all possible states of quantum many-particle systems in comparison with the approach formulated above.

In what follows we will use such abridged notations: $Y \equiv (1,\ldots,n)$ and $Y_{\mathrm{P}} \equiv (X_1,\ldots,X_{|\mathrm{P}|})$ is a set whose elements are $|\mathrm{P}|$ mutually disjoint subsets $X_i \subset Y$ of the partition $\mathrm{P} : Y = \bigcup_{i=1}^{|\mathrm{P}|} X_i$. The $|\mathrm{P}|th$-order cumulant (1.9) in this case we denote by $\mathfrak{A}_{|\mathrm{P}|}(t, Y_{\mathrm{P}})$.

A solution of an initial-value problem to the von Neumann hierarchy is defined by a one-parametric family of nonlinear operators [15] constructed in [23] with the following properties.

Theorem 2.1. *If $f_n \in \mathfrak{L}^1(\mathcal{H}_n)$, $n \geq 1$, then the one-parametric family of nonlinear operators*

$$\mathbb{R}^1 \ni t \mapsto \big(\mathfrak{A}_t(f)\big)_n(Y) := \sum_{\mathrm{P}:\, Y=\bigcup_i X_i} \mathfrak{A}_{|\mathrm{P}|}(t, Y_{\mathrm{P}}) \prod_{X_i \subset \mathrm{P}} f_{|X_i|}(X_i) \tag{2.1}$$

is a C_0-group. On the subspace $\mathfrak{L}^1_0(\mathcal{H}_n) \subset \mathfrak{L}^1(\mathcal{H}_n)$ the infinitesimal generator $\mathfrak{N}$ of group (2.1) *is defined by the operator*

$$\begin{aligned}\big(\mathfrak{N}(f)\big)_n(Y) := {}& -\mathcal{N}_n(Y) f_n(Y) + \\ & \sum_{\substack{\mathrm{P}:\, Y = \bigcup_i X_i,\\ |\mathrm{P}| > 1}} \; \sum_{\substack{Z_1 \subset X_1,\\ Z_1 \neq \emptyset}} \cdots \sum_{\substack{Z_{|\mathrm{P}|} \subset X_{|\mathrm{P}|},\\ Z_{|\mathrm{P}|} \neq \emptyset}} \Big(-\mathcal{N}_{\mathrm{int}}^{\big(\sum_{r=1}^{|\mathrm{P}|} |Z_r|\big)} \big(Z_1,\ldots,Z_{|\mathrm{P}|}\big)\Big) \prod_{X_i \subset \mathrm{P}} f_{|X_i|}(X_i),\end{aligned} \tag{2.2}$$

where the notations are similar to that in (1.8) *and the operator $\mathcal{N}_{\mathrm{int}}^{(n)}$ is defined by formula* (1.11).

Proof. Mapping (2.1) is defined for $f_n \in \mathfrak{L}^1(\mathcal{H}_n)$, $n \geq 1$, and the following inequality holds

$$\big\| \big(\mathfrak{A}_t(f)\big)_n \big\|_{\mathfrak{L}^1(\mathcal{H}_n)} \leq n! e^{2n+1} c^n, \tag{2.3}$$

where $c := \max\limits_{\mathrm{P}:\, Y=\bigcup_i X_i} \|f_{|X_i|}(X_i)\|_{\mathfrak{L}^1(\mathcal{H}_{|X_i|})}$. Indeed, since for $f_n \in \mathfrak{L}^1(\mathcal{H}_n)$ the equality holds [23]

$$\mathrm{Tr}_{1,\ldots,n} |\mathcal{G}_n(-t) f_n| = \|f_n\|_{\mathfrak{L}^1(\mathcal{H}_n)},$$

we have

$$\begin{aligned}\big\| \big(\mathfrak{A}_t(f)\big)_n \big\|_{\mathfrak{L}^1(\mathcal{H}_n)} \leq \sum_{\mathrm{P}:\, Y=\bigcup_i X_i} \; \sum_{\mathrm{P}':\, Y_{\mathrm{P}}=\bigcup_k Z_k} (|\mathrm{P}'| - 1)! \prod_{X_i \subset \mathrm{P}} \|f_{|X_i|}\|_{\mathfrak{L}^1(\mathcal{H}_{|X_i|})} \leq \\ \sum_{\mathrm{P}:\, Y=\bigcup_i X_i} c^{|\mathrm{P}|} \sum_{k=1}^{|\mathrm{P}|} s(|\mathrm{P}|, k)(k-1)! \leq \sum_{\mathrm{P}:\, Y=\bigcup_i X_i} c^{|\mathrm{P}|} \sum_{k=1}^{|\mathrm{P}|} k^{|\mathrm{P}|-1} \leq n! e^{2n+1} c^n,\end{aligned}$$

where $\mathrm{s}(|\mathrm{P}|, k)$ are the Stirling numbers of the second kind. That is, $\big(\mathfrak{A}_t(f)\big)_n \in \mathfrak{L}^1(\mathcal{H}_n)$ for arbitrary $t \in \mathbb{R}^1$ and $n \geq 1$.

The group property of a one-parametric family of nonlinear operators (2.1), i.e. $\mathfrak{A}_{t_1}\big(\mathfrak{A}_{t_2}(f)\big) = \mathfrak{A}_{t_2}\big(\mathfrak{A}_{t_1}(f)\big) = \mathfrak{A}_{t_1+t_2}(f)$, was proved in [23].

The strong continuity property of the group $\{\mathfrak{A}_t\}_{t\in\mathbb{R}}$ over the parameter $t\in\mathbb{R}^1$ is a consequence of the strong continuity of group (1.1) of the von Neumann equation [10]. Indeed, according to identity (1.10) the following equality holds:

$$\sum_{\mathrm{P}:\,Y=\bigcup_i X_i}\ \sum_{\mathrm{P}':\,Y_{\mathrm{P}}=\bigcup_k Z_k}(-1)^{|\mathrm{P}'|-1}(|\mathrm{P}'|-1)!\prod_{X_i\subset\mathrm{P}} f_{|X_i|}(X_i)=f_n(Y),$$

Therefore, for $f_n\in\mathfrak{L}^1_0(\mathcal{H}_n)\subset\mathfrak{L}^1(\mathcal{H}_n)$, $n\geq 1$, we have

$$\begin{aligned}&\lim_{t\rightarrow 0}\big\|\big(\mathfrak{A}_t(f)\big)_n(Y)-f_n(Y)\big\|_{\mathfrak{L}^1(\mathcal{H}_n)}\leq\\ &\sum_{\mathrm{P}:\,Y=\bigcup_i X_i}\ \sum_{\mathrm{P}':\,Y_{\mathrm{P}}=\bigcup_k Z_k}(|\mathrm{P}'|-1)!\,\lim_{t\rightarrow 0}\big\|\big(\prod_{Z_k\subset\mathrm{P}'}\mathcal{G}_{|Z_k|}(-t,Z_k)-I\big)\prod_{X_i\subset\mathrm{P}} f_{|X_i|}(X_i)\big\|_{\mathfrak{L}^1(\mathcal{H}_n)}.\end{aligned}$$

In view of the the fact that group $\{\mathcal{G}_n(-t)\}_{t\in\mathbb{R}}$ (1.1) is a strong continuous group, which implies that, for mutually disjoint subsets $X_i\subset Y$, if $f_n\in\mathfrak{L}^1_0(\mathcal{H}_n)\subset\mathfrak{L}^1(\mathcal{H}_n)$ in the sense of the norm convergence $\mathfrak{L}^1(\mathcal{H}_n)$ there exists the limit

$$\lim_{t\rightarrow 0}\big(\prod_{Z_k\subset\mathrm{P}'}\mathcal{G}_{|Z_k|}(-t,Z_k)f_n-f_n\big)=0.$$

Thus if $f\in\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$ we finally obtain

$$\lim_{t\rightarrow 0}\big\|\big(\mathfrak{A}_t(f)\big)_n-f_n\big\|_{\mathfrak{L}^1(\mathcal{H}_n)}=0.$$

We now construct the infinitesimal generator $\mathfrak{N}$ of group (2.1). Taking into account that for $f_n\in\mathfrak{L}^1_0(\mathcal{H}_n)$ equality (1.4) holds, we differentiate the $|\mathrm{P}|th$-order cumulant $\mathfrak{A}_{|\mathrm{P}|}(t,Y_{\mathrm{P}})$ for all $\psi_n\in\mathcal{D}(H_n)\subset\mathcal{H}_n$ in the sense of the point-by-point convergence. According to equality (1.12) for $|\mathrm{P}|\geq 2$ we derive

$$\begin{aligned}\lim_{t\rightarrow 0}\frac{1}{t}\mathfrak{A}_{|\mathrm{P}|}(t,Y_{\mathrm{P}})f_n\psi_n&=\sum_{\mathrm{P}':\,Y_{\mathrm{P}}=\bigcup_k Z_k}(-1)^{|\mathrm{P}'|-1}(|\mathrm{P}'|-1)!\sum_{Z_k\subset\mathrm{P}'}(-\mathcal{N}_{|Z_k|}(Z_k))f_n\psi_n\\ &=\sum_{\substack{Z_1\subset X_1,\\ Z_1\neq\emptyset}}\ldots\sum_{\substack{Z_{|\mathrm{P}|}\subset X_{|\mathrm{P}|},\\ Z_{|\mathrm{P}|}\neq\emptyset}}\Big(-\mathcal{N}_{\mathrm{int}}^{\big(\sum\limits_{r=1}^{|\mathrm{P}|}|Z_r|\big)}\big(Z_1,\ldots,Z_{|\mathrm{P}|}\big)\Big)f_n\psi_n,\end{aligned}\tag{2.4}$$

where $\sum_{Z_j\subset X_j}$ is a sum over all subsets $Z_j\subset X_j$ of the set X_j. Then in view of equality (2.4) for group (2.1) we obtain

$$\begin{aligned}&\lim_{t\rightarrow 0}\frac{1}{t}\Big(\big(\mathfrak{A}_t(f)\big)_n-f_n\Big)\psi_n=\lim_{t\rightarrow 0}\frac{1}{t}\Big(\sum_{\mathrm{P}:\,Y=\bigcup_i X_i}\mathfrak{A}_{|\mathrm{P}|}(t,Y_{\mathrm{P}})\prod_{X_i\subset\mathrm{P}}f_{|X_i|}(X_i)-f_n(Y)\Big)\psi_n\\ &=\lim_{t\rightarrow 0}\frac{1}{t}\big(\mathfrak{A}_1(t,Y)f_n-f_n\big)\psi_n+\sum_{\substack{\mathrm{P}\,:\,Y=\bigcup_i X_i,\\ |\mathrm{P}|>1}}\lim_{t\rightarrow 0}\frac{1}{t}\mathfrak{A}_{|\mathrm{P}|}(t,Y_{\mathrm{P}})\prod_{X_i\subset\mathrm{P}}f_{|X_i|}(X_i)\psi_n\\ &=\big(-\mathcal{N}_n f_n\big)(Y)\psi_n\\ &+\sum_{\substack{\mathrm{P}\,:\,Y=\bigcup_i X_i,\\ |\mathrm{P}|>1}}\ \sum_{\substack{Z_1\subset X_1,\\ Z_1\neq\emptyset}}\ldots\sum_{\substack{Z_{|\mathrm{P}|}\subset X_{|\mathrm{P}|},\\ Z_{|\mathrm{P}|}\neq\emptyset}}\Big(-\mathcal{N}_{\mathrm{int}}^{\big(\sum\limits_{r=1}^{|\mathrm{P}|}|Z_r|\big)}\big(Z_1,\ldots,Z_{|\mathrm{P}|}\big)\Big)\prod_{X_i\subset\mathrm{P}}f_{|X_i|}(X_i)\psi_n.\end{aligned}$$

Thus for $f \in \mathfrak{L}^1_0(\mathcal{F}_\mathcal{H}) \subset \mathcal{D}(\mathfrak{N}) \subset \mathfrak{L}^1(\mathcal{F}_\mathcal{H})$ in the sense of the norm convergence $\mathfrak{L}^1(\mathcal{H}_n)$ we finally have

$$\lim_{t\rightarrow 0}\big\|\frac{1}{t}\big(\big(\mathfrak{A}_t(f)\big)_n - f_n\big) - \big(\mathfrak{N}(f)\big)_n\big\|_{\mathfrak{L}^1(\mathcal{H}_n)} = 0. \qquad \square$$

We give an example which illustrates the structure of expansion (2.1). For the correlation operators $f = \big(0, f_1(1), 0, \ldots\big)$ that is interpreted as satisfying the "chaos" property [1], we have

$$\big(\mathfrak{A}_t(f)\big)_n = \mathfrak{A}_n(t, 1, \ldots, n)\prod_{i=1}^n f_1(i), \quad n \geq 1,$$

i.e. if at the initial instant there are no correlations in a system, the correlations generated by the dynamics of a system are completely governed by cumulants of groups (1.1).

We now consider the structure of infinitesimal generator (2.2) for a two-body interaction potential

$$\big(\mathfrak{N}(f)\big)_n(Y) = -\mathcal{N}_n(Y)f_n(Y) + \sum_{\mathrm{P}:\, Y = X_1\bigcup X_2}\ \sum_{i_1\in\{X_1\}}\sum_{i_2\in\{X_2\}}\big(-\mathcal{N}^{(2)}_{\mathrm{int}}(i_1,i_2)\big)f_{|X_1|}(X_1)f_{|X_2|}(X_2),$$

where the symbol $\sum_{\mathrm{P}:\, Y=X_1\bigcup X_2}$ means summation over all partitions of the set Y into two nonempty parts X_1 and X_2 and the operator $\mathcal{N}^{(2)}_{\mathrm{int}}$ is defined by formula (1.11). For classical systems this generator is an equivalent notation of the generator of the Liouville hierarchy [20] formulated in [21].

3 Group of Operators for the Quantum BBGKY Hierarchy

The evolution of all possible states both finitely and infinitely many quantum particles is described by the initial-value problem to the BBGKY hierarchy for marginal density operators [11],[24]. For finitely many particles this hierarchy of equations is an equivalent to the von Neumann equation.

We will use notations from the previous section. Since $Y_\mathrm{P} \equiv (X_1, \ldots, X_{|\mathrm{P}|})$ then Y_1 is the set consisting of one element of the partition P ($|\mathrm{P}| = 1$) of the set $Y \equiv (1, \ldots, s)$. In this case for $n \geq 0$ the $(1+n)th$-order cumulant of operators (1.1) is defined by the formula

$$\mathfrak{A}_{1+n}(t, Y_1, X\backslash Y) := \sum_{\mathrm{P}:\,\{Y_1, X\backslash Y\} = \bigcup_i X_i} (-1)^{|\mathrm{P}|-1}(|\mathrm{P}|-1)!\prod_{X_i\subset \mathrm{P}}\mathcal{G}_{|X_i|}(-t, X_i), \tag{3.1}$$

where $\sum_\mathrm{P}$ is the sum over all possible partitions P of the set $\{Y_1, X\setminus Y\} = \{Y_1, s+1, \ldots, s+n\}$ into $|\mathrm{P}|$ nonempty mutually disjoint subsets $X_i \subset \{Y_1, X\setminus Y\}$.

On the space $\mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$ a solution of the initial-value problem to the BBGKY hierarchy is defined by a one-parametric mapping [24] with the following properties.

Theorem 3.1. *If $f \in \mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$ and $\alpha > e$, then the one-parametric mapping*

$$\mathbb{R}^1 \ni t \mapsto (U(t)f)_s(Y) := \sum_{n=0}^\infty \frac{1}{n!}\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}\mathfrak{A}_{1+n}(t, Y_1, X\backslash Y)f_{s+n}(X) \tag{3.2}$$

is a C_0-group. On the subspace $\mathfrak{L}^1_{\alpha,0} \subset \mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$ the infinitesimal generator $\mathfrak{B} = \bigoplus_{n=0}^\infty \mathfrak{B}_n$ of group (3.2) *is defined by the operator* ($s \geq 1$)

$$\begin{aligned}(\mathfrak{B}f)_s(Y) &:= \\ &-\mathcal{N}_s(Y)f_s(Y) + \sum_{k=1}^s\frac{1}{k!}\sum_{i_1\neq\ldots\neq i_k=1}^s\ \sum_{n=1}^\infty\frac{1}{n!}\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}\big(-\mathcal{N}^{(k+n)}_{\mathrm{int}}\big)(i_1,\ldots,i_k, X\backslash Y)f_{s+n}(X),\end{aligned} \tag{3.3}$$

where on $\mathfrak{L}^1_0(\mathcal{H}_{s+n}) \subset \mathfrak{L}^1(\mathcal{H}_{s+n})$ the operator $\mathcal{N}^{(k+n)}_{\mathrm{int}}$ is defined by formula (1.11).

Proof. If $f \in \mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$ mapping (3.2) is defined provided that $\alpha > e$ and the following estimate holds [24]

$$\|U(t)f\|_{\mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})} \leq c_\alpha \|f\|_{\mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})},$$

where $c_\alpha = e^2(1-\frac{e}{\alpha})^{-1}$ is a constant. Similar to (2.3) this estimate comes out from the inequality for cumulant (3.1)

$$\|\mathfrak{A}_{1+n}(t) f_{s+n}\|_{\mathfrak{L}^1(\mathcal{H}_{s+n})} \leq n! e^{n+2} \|f_{s+n}\|_{\mathfrak{L}^1(\mathcal{H}_{s+n})}.$$

The strong continuity property of the group $U(t)$ over the parameter $t \in \mathbb{R}^1$ is a consequence of the strong continuity of group (1.1) of the von Neumann equation.

We now construct an infinitesimal generator of group (3.2). Taking into account that for $f_n \in \mathfrak{L}^1_0(\mathcal{H}_n)$ equality (1.4) holds, we differentiate the expression of cumulant (3.1) in the sense of the point-by-point convergence. According to equality (2.4) for $n \geq 1$, we derive

$$\begin{aligned}&\lim_{t\rightarrow 0}\frac{1}{t}\mathfrak{A}_{1+n}(t, Y_1, X\backslash Y) f_{s+n}\psi_{s+n} = \sum_{\substack{Z\subset Y,\\ Z\neq\emptyset}} \big(-\mathcal{N}_{\mathrm{int}}^{(|Z|+n)}\big)(Z, s+1, \ldots, s+n) f_{s+n}\psi_{s+n} = \\ &\sum_{k=1}^{|Y|}\frac{1}{k!}\sum_{i_1\neq\ldots\neq i_k=1}^{|Y|}\big(-\mathcal{N}_{\mathrm{int}}^{(k+n)}\big)(i_1,\ldots,i_k, X\backslash Y) f_{s+n}\psi_{s+n},\end{aligned} \tag{3.4}$$

where $\sum_{Z\subset X}$ is a sum over all subsets $Z \subset X$ of the set X. Then taking into account formula (1.4) for $n = 1$ and equality (3.4) for group (3.2) we obtain

$$\begin{aligned}&\lim_{t\rightarrow 0}\frac{1}{t}\big(\big(U(t)f\big)_s - f_s\big)\psi_s = \\ &\lim_{t\rightarrow 0}\frac{1}{t}\big(\mathfrak{A}_1(t,Y) f_s - f_s\big)\psi_s + \sum_{n=1}^{\infty}\frac{1}{n!}\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}\lim_{t\rightarrow 0}\frac{1}{t}\mathfrak{A}_{1+n}(t, Y_1, X\backslash Y) f_{s+n}\psi_s = \\ &-\mathcal{N}_s f_s\psi_s + \sum_{n=1}^{\infty}\frac{1}{n!}\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}\sum_{k=1}^{s}\frac{1}{k!}\sum_{i_1\neq\ldots\neq i_k=1}^{s}\big(-\mathcal{N}_{\mathrm{int}}^{(k+n)}\big)(i_1,\ldots,i_k, X\backslash Y)\psi_s.\end{aligned}$$

Thus, if $\mathfrak{L}^1_{\alpha,0} \subset \mathcal{D}(\mathfrak{B}) \subset \mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$ we finally have in the sense of the norm convergence

$$\lim_{t\rightarrow 0}\big\|\frac{1}{t}\big(U(t)f - f\big) - \mathfrak{B}f\big\|_{\mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})} = 0,$$

where the operator $\mathfrak{B}$ on $\mathfrak{L}^1_{\alpha,0}$ is given by formula (3.3). □

We now give an example which illustrates the structure of infinitesimal generator (3.3). In the case of two-body interaction potential (1.3) operator (3.3) has the form ($s \geq 1$)

$$(\mathfrak{B}f)_s(Y) = -\mathcal{N}_s(Y) f_s(Y) + \sum_{i=1}^{s}\mathrm{Tr}_{\mathrm{s+1}}\big(-\mathcal{N}_{\mathrm{int}}^{(2)}\big)(i, s+1) f_{s+1}(Y, s+1), \tag{3.5}$$

where the operator $\mathcal{N}_{int}^{(2)}$ is defined on $\mathfrak{L}^1_0(\mathcal{H}_{s+1}) \subset \mathfrak{L}^1(\mathcal{H}_{s+1})$ by formula (1.11) for $n = 2$. For $\mathcal{H} = L^2(\mathbb{R}^3)$ in the framework of kernels of operators f_s (s-particle density matrix or marginal distributions [11]) operator (3.5) takes a canonical form of a generator of the quantum BBGKY hierarchy [1],[24]

$$
\begin{aligned}
&(\mathfrak{B}f)_s(q_1,\ldots,q_s;q_1^{'},\ldots,q_s^{'}) = \\
&-\frac{i}{\hbar}\Big(-\frac{\hbar^2}{2}\sum_{i=1}^{s}(\Delta_{q_i}-\Delta_{q_i^{'}})+\sum_{i<j=1}^{s}\big(\Phi^{(2)}(q_i-q_j)-\Phi^{(2)}(q_i^{'}-q_j^{'})\big)\Big)f_s(q_1,\ldots,q_s;q_1^{'},\ldots,q_s^{'})- \\
&\frac{i}{\hbar}\sum_{i=1}^{s}\int dq_{s+1}\big(\Phi^{(2)}(q_i-q_{s+1})-\Phi^{(2)}(q_i^{'}-q_{s+1})\big)f_{s+1}(q_1,\ldots q_s,q_{s+1};q_1^{'},\ldots,q_s^{'},q_{s+1}).
\end{aligned}
$$

In [11],[24] for quantum and in the book [1] for classical systems of particles with a two-body interaction potential, an equivalent representation for group (3.2) was used, namely for the case under consideration the group $U(t)$ has the following representation

$$
U(t)=\mathcal{G}(-t)+\sum_{n=1}^{\infty}\frac{1}{n!}\big[\underbrace{\mathfrak{a},\ldots,\big[\mathfrak{a}}_{\text{n-times}},\mathcal{G}(-t)\big]\ldots\big]=e^{\mathfrak{a}}\mathcal{G}(-t)e^{-\mathfrak{a}}, \tag{3.6}
$$

where $\big[\,.\,,\,.\,\big]$ is a commutator, the operator $\mathfrak{a}$ (an analog of the annihilation operator) is defined on the space $\mathfrak{L}^1_{\alpha}(\mathcal{F}_{\mathcal{H}})$ by the formula

$$
\big(\mathfrak{a}f\big)_s(Y):=\mathrm{Tr}_{\mathrm{s+1}}f_{s+1}(Y,s+1), \tag{3.7}
$$

the operators $e^{\pm\mathfrak{a}}$ are defined on the space $\mathfrak{L}^1_{\alpha}(\mathcal{F}_{\mathcal{H}})$ by the expansions

$$
\big(e^{\pm\mathfrak{a}}f\big)_s(Y)=\sum_{n=0}^{\infty}\frac{(\pm1)^n}{n!}\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}f_{s+n}(Y,s+1,\ldots,s+n),\quad s\geq1.
$$

and the group of operators $\mathcal{G}(-t)$ is defined by expression (1.1). Representation (3.6) is true in consequence of definition (3.7) of the operator $\mathfrak{a}$ and the validity of an equality for every $s\geq1$

$$
\big(\big[\underbrace{\mathfrak{a},\ldots,\big[\mathfrak{a}}_{\text{n-times}},\mathcal{G}(-t)\big]\ldots\big]f\big)_s(Y)=\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}\mathfrak{A}_{1+n}(t,Y_1,X\backslash Y)f_{s+n}(X),
$$

where the notations are similar to that in (3.1). For example, since for an isometric group $\mathcal{G}(-t)$ and trace class operators f the following equality holds

$$
\mathrm{Tr}_{\mathrm{s+1}}\mathcal{G}_1(-t,s+1)f_{s+1}(Y,s+1)=\mathrm{Tr}_{\mathrm{s+1}}f_{s+1}(Y,s+1). \tag{3.8}
$$

then if $n=1$ we have

$$
\begin{aligned}
&\big(\big[\mathfrak{a},\mathcal{G}(-t)\big]f\big)_s(Y)=\mathrm{Tr}_{\mathrm{s+1}}\big(\mathcal{G}_{s+1}(-t,Y,s+1)f_{s+1}(Y,s+1)-\mathcal{G}_s(-t,Y)f_{s+1}(Y,s+1)\big)= \\
&\mathrm{Tr}_{\mathrm{s+1}}\mathfrak{A}_2(t,Y_1,s+1)f_{s+1}(Y,s+1).
\end{aligned}
$$

We note that cluster expansions (1.8) can be put into basis of all possible representations of a group of operators for the quantum BBGKY hierarchy. In fact representation (3.6) we obtain solving recurrence relations (1.8) with respect to the $1st$-order cumulants for the separation terms which are independent from the variable Y

$$
\mathfrak{A}_{1+n}\big(t,Y_1,X\backslash Y\big)=\sum_{Z\subset X\backslash Y}\mathfrak{A}_1(t,Y\cup Z)\sum_{\mathrm{P}:\,(X\backslash Y)\backslash Z=\bigcup_i X_i}(-1)^{|\mathrm{P}|}\,|\mathrm{P}|!\,\prod\nolimits_{i=1}^{|\mathrm{P}|}\mathfrak{A}_1(t,X_i),
$$

where $\sum_{Z\subset X\backslash Y}$ is a sum over all subsets $Z\subset X\backslash Y$ of the set $X\backslash Y$. Then taking into account equality (3.8) and

$$\sum_{\mathrm{P}\colon (X\backslash Y)\backslash Z=\bigcup_i X_i}(-1)^{|\mathrm{P}|}\,|\mathrm{P}|!=(-1)^{|(X\backslash Y)\backslash Z|},\tag{3.9}$$

for the group $U(t)$ we have

$$\big(U(t)f\big)_s(Y)=\sum_{n=0}^{\infty}\frac{1}{n!}\mathrm{Tr}_{s+1,\ldots,s+n}\sum_{Z\subset X\backslash Y}(-1)^{|(X\backslash Y)\backslash Z|}\mathcal{G}_{|Y\cup Z|}(-t,Y\cup Z)f_{|X|}(X).$$

Thus as a result of the symmetry property and definition (3.7) of the operator $\mathfrak{a}$ we derive (3.6)

$$U(t)=\sum_{n=0}^{\infty}\frac{1}{n!}\sum_{k=0}^{n}(-1)^k\frac{n!}{k!(n-k)!}\mathfrak{a}^{n-k}\mathcal{G}(-t)\mathfrak{a}^k=e^{\mathfrak{a}}\mathcal{G}(-t)e^{-\mathfrak{a}}.$$

We can obtain one more representation of the group $U(t)$ solving recurrence relations (1.8) with respect to the 1*st*-order and 2*nd*-order cumulants. In fact if $n\geq 1$ we get

$$\mathfrak{A}_{1+n}(t,Y_1,X\backslash Y)=\sum_{\substack{Z\subset X\backslash Y,\\ Z\neq\emptyset}}\mathfrak{A}_2(t,Y,Z)\sum_{\mathrm{P}\colon (X\backslash Y)\backslash Z=\bigcup_i X_i}(-1)^{|\mathrm{P}|}\,|\mathrm{P}|!\prod\nolimits_{i=1}^{|\mathrm{P}|}\mathfrak{A}_1(t,X_i).$$

Then taking into account equalities (3.8) and (3.9) we derive

$$\big(U(t)f\big)_s(Y)=\mathfrak{A}_1(t,Y)f_s(Y)+\sum_{n=1}^{\infty}\frac{1}{n!}\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}\sum_{\substack{Z\subset X\backslash Y,\\ Z\neq\emptyset}}(-1)^{|(X\backslash Y)\backslash Z|}\mathfrak{A}_2(t,Y,Z)f_{s+n}(X).$$

An infinitesimal generator of a group of operators (3.6) has the following form

$$\mathfrak{B}=-\mathcal{N}+\sum_{n=1}^{\infty}\frac{1}{n!}\big[\underbrace{\mathfrak{a},\ldots,\big[\mathfrak{a}}_{\text{n-times}},(-\mathcal{N})\big]\ldots\big]=e^{\mathfrak{a}}(-\mathcal{N})e^{-\mathfrak{a}}.\tag{3.10}$$

Representation (3.10) is true in consequence of definition (3.7) of the operator $\mathfrak{a}$ and the validity of equalities

$$\begin{aligned}&\big(\big[\underbrace{\mathfrak{a},\ldots,\big[\mathfrak{a}}_{\text{n-times}},(-\mathcal{N})\big]\ldots\big]f\big)_s(Y)=\sum_{k=0}^{n}(-1)^k\frac{n!}{k!(n-k)!}\big(\mathfrak{a}^{n-k}(-\mathcal{N})\mathfrak{a}^kf\big)_s(Y)=\\&\sum_{k=0}^{n}(-1)^k\sum_{i_1<\ldots<i_{n-k}=s+1}^{s+n}\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}\big(-\mathcal{N}_{s+n-k}(Y,i_1,\ldots,i_{n-k})\big)f_{s+n}(X)=\\&\sum_{k=1}^{s}\frac{1}{k!}\sum_{i_1\neq\ldots\neq i_k=1}^{s}\mathrm{Tr}_{\mathrm{s+1,\ldots,s+n}}\big(-\mathcal{N}_{\mathrm{int}}^{(k+n)}\big)(i_1,\ldots,i_k,X\backslash Y)f_{s+n}(X).\end{aligned}$$

For a system of particles interacting through a two-body potential this equality reduces to the following one

$$\big([\mathcal{N},\mathfrak{a}]f\big)_s(Y)=\sum_{i=1}^{s}\mathrm{Tr}_{\mathrm{s+1}}\big(-\mathcal{N}_{\mathrm{int}}^{(2)}\big)(i,s+1)f_{s+1}(Y,s+1),$$

that is true in view of the equality: $\mathrm{Tr}_{\mathrm{s+1}}\mathcal{N}_1(s+1)f_{s+1}(Y,s+1)=0$.

4 Group of Operators for the Quantum Dual BBGKY Hierarchy

The evolution of marginal observables of both finitely and infinitely many quantum particles is described by the initial-value problem to the dual BBGKY hierarchy. This hierarchy of equations is dual to the quantum BBGKY hierarchy in the sense of bilinear form (1.5) and for finitely many particles one is an equivalent to the Heisenberg equation (the dual von Neumann equation). For systems of classical particles the dual BBGKY hierarchy was examined in [1],[25],[26].

In this section we will use such abridged notations: $Y \equiv (1,\ldots,s)$, $X \equiv Y\backslash\{j_1,\ldots,j_{s-n}\}$. According to notations of section 2, the set $(Y\backslash X)_1$ consists of one element $Y\backslash X = (j_1,\ldots,j_{s-n})$, i.e. the set $(j_1,\ldots,j_{s-n})$ is connected subset of the partition P ($|\mathrm{P}| = 1$). In the case under consideration the dual cumulants $\mathfrak{A}^{+}_{1+n}(t),\ n\geq 0$, of groups (1.6) are defined by the formula

$$\mathfrak{A}^{+}_{1+n}\bigl(t,(Y\backslash X)_1,X\bigr) := \sum_{\mathrm{P}:\,\{(Y\backslash X)_1,X\}=\bigcup_i X_i} (-1)^{|\mathrm{P}|-1}(|\mathrm{P}|-1)! \prod_{X_i\subset \mathrm{P}} \mathcal{G}_{|X_i|}(t,X_i), \tag{4.1}$$

where $\sum_{\mathrm{P}}$ is the sum over all possible partitions P of the set $\{(Y\backslash X)_1, j_1,\ldots,j_{s-n}\}$ into $|\mathrm{P}|$ nonempty mutually disjoint subsets $X_i \subset \{(Y\backslash X)_1, X\}$.

On the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ a solution of the initial-value problem to the dual BBGKY hierarchy is defined by a one-parametric mapping (the adjoint mapping to (3.2) in the sense of bilinear form (1.5)) with the following properties.

Theorem 4.1. *If $g\in\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ and $\gamma<e^{-1}$, then the one-parametric mapping*

$$\begin{aligned}&\mathbb{R}^1 \ni t \mapsto \bigl(U^+(t)g\bigr)_s(Y) := \\ &\sum_{n=0}^{s}\frac{1}{(s-n)!}\sum_{j_1\neq\ldots\neq j_{s-n}=1}^{s} \mathfrak{A}^{+}_{1+n}\bigl(t,(Y\backslash X)_1,X\bigr)\, g_{s-n}(Y\backslash X), \quad s\geq 1\end{aligned} \tag{4.2}$$

is a C_0^*-group. The infinitesimal generator $\mathfrak{B}^+ = \bigoplus_{n=0}^{\infty}\mathfrak{B}^+_n$ of this group of operators is a closed operator for the $*$-weak topology and on the domain of the definition $\mathcal{D}(\mathfrak{B}^+)\subset\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ which is the everywhere dense set for the $*$-weak topology of the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ it is defined by the operator

$$\begin{aligned}&(\mathfrak{B}^+g)_s(Y) := \mathcal{N}_s(Y)g_s(Y) + \\ &\sum_{n=1}^{s}\frac{1}{n!}\sum_{k=n+1}^{s}\frac{1}{(k-n)!}\sum_{j_1\neq\ldots\neq j_k=1}^{s} \mathcal{N}^{(k)}_{\mathrm{int}}(j_1,\ldots,j_k)g_{s-n}(Y\backslash\{j_1,\ldots,j_n\}),\end{aligned} \tag{4.3}$$

where the operator $\mathcal{N}^{(k)}_{\mathrm{int}}$ is given by formula (1.11).

Proof. If $g\in\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ mapping (4.2) is defined provided that $\gamma<e^{-1}$ and the following estimate holds

$$\bigl\|U^+(t)g\bigr\|_{\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})} \leq e^2(1-\gamma e)^{-1}\|g\|_{\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})}.$$

Similar to (2.3) this estimate comes out from the inequality

$$\begin{aligned}&\bigl\|(U^+(t)g)_s\bigr\|_{\mathfrak{L}(\mathcal{H}_s)} \leq \sum_{n=0}^{s}\frac{1}{(s-n)!}\sum_{j_1\neq\ldots\neq j_{s-n}=1}^{s}\ \sum_{\mathrm{P}:\{Y\backslash X)_1,X\}=\bigcup_i X_i}(|\mathrm{P}|-1)!\|g_{s-n}\|_{\mathfrak{L}(\mathcal{H}_{s-n})} \leq \\ &\sum_{n=0}^{s}\|g_{s-n}\|_{\mathfrak{L}(\mathcal{H}_{s-n})}\frac{s!}{n!(s-n)!}\sum_{k=1}^{n+1}s(n+1,k)(k-1)!,\end{aligned}$$

where $s(n+1,k)$ is the Stirling numbers of the second kind.

On the space $\mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$ the $*$-weak continuity property of the group $U^{+}(t)$ over the parameter $t\in\mathbb{R}^{1}$ is a consequence of the $*$-weak continuity of group (1.6) of the Heisenberg equation [10].

To construct an infinitesimal generator of the group $\{U^{+}(t)\}_{t\in\mathbb{R}}$ we firstly differentiate the *nth*-term of expansion (4.2) in the sense of the point-by-point convergence of the space $\mathfrak{L}_{\gamma}$. If $g\in\mathcal{D}(\mathcal{N})\subset\mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$ similar to equality (1.12) for $(1+n)th$-order dual cumulant (4.1), $n\geq 1$, we derive

$$\lim_{t\rightarrow 0}\frac{1}{t}\mathfrak{A}^{+}_{1+n}\big(t,(Y\backslash X)_{1},X\big)g_{s-n}(Y\backslash X)\psi_{s}=\sum_{\substack{Z\subset Y\backslash X,\\ Z\neq\emptyset}}\mathcal{N}^{(|Z|+n)}_{\mathrm{int}}(Z,X)g_{s-n}(Y\backslash X)\psi_{s}=$$
$$\sum_{k=1}^{s-n}\frac{1}{k!}\sum_{i_{1}\neq\ldots\neq i_{k}\in\{j_{1},\ldots,j_{s-n}\}}\mathcal{N}^{(k+n)}_{\mathrm{int}}(i_{1},\ldots,i_{k},X)g_{s-n}(Y\backslash X)\psi_{s}. \tag{4.4}$$

Then according to equalities (1.7) and (4.4) for group (4.2) we obtain

$$\lim_{t\rightarrow 0}\frac{1}{t}\big(\big(U^{+}(t)g\big)_{s}-g_{s}\big)\psi_{s}=\lim_{t\rightarrow 0}\frac{1}{t}\big(\mathfrak{A}^{+}_{1}(t)g_{s}-g_{s}\big)\psi_{s}+$$
$$\sum_{n=1}^{s}\frac{1}{(s-n)!}\sum_{j_{1}\neq\ldots\neq j_{s-n}=1}^{s}\lim_{t\rightarrow 0}\frac{1}{t}\mathfrak{A}^{+}_{1+n}\big(t,(Y\backslash X)_{1},X\big)g_{s-n}(Y\backslash X)\psi_{s}=$$
$$\mathcal{N}_{s}g_{s}\psi_{s}+\sum_{n=1}^{s}\frac{1}{n!}\sum_{k=n+1}^{s}\frac{1}{(k-n)!}\sum_{j_{1}\neq\ldots\neq j_{k}=1}^{s}\mathcal{N}^{(k)}_{\mathrm{int}}(j_{1},\ldots,j_{k})g_{s-n}(Y\backslash\{j_{1},\ldots,j_{n}\})\psi_{s},$$

where we used the identity

$$\sum_{n=0}^{s}\frac{1}{n!}\sum_{j_{1}\neq\ldots\neq j_{n}=1}^{s}g_{s-n}(Y\backslash\{j_{1},\ldots,j_{n}\})=\sum_{n=0}^{s}\frac{1}{n!}\sum_{j_{1}\neq\ldots\neq j_{n}=1}^{s}g_{n}(j_{1},\ldots,j_{n}) \tag{4.5}$$

which is valid in view of the Maxwell-Boltzmann statistics symmetry property.

Thus if $g\in\mathcal{D}(\mathfrak{B}^{+})\subset\mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$ in the sense of the $*$-weak convergence of the space $\mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$ we finally have

$$\mathrm{w}^{*}\!-\lim_{t\rightarrow 0}\Big(\frac{1}{t}\big(U^{+}(t)g-g\big)-\mathfrak{B}^{+}g\Big)=0,$$

where the generator $\mathfrak{B}^{+}=\bigoplus_{n=0}^{\infty}\mathfrak{B}^{+}_{n}$ of group (4.2) is given by formula (4.3) (the dual operator to generator (3.3)). □

We now give examples of expansion (4.2) and infinitesimal generator (4.3). The sequence $g=\big(0,g_{1}(1),0,\ldots\big)$ corresponds to the additive-type observable [25] and in this case expansion (4.2) for the group $U^{+}(t)$ get the form

$$(U^{+}(t)g)_{s}(Y)=\mathfrak{A}^{+}_{s}(t,1,\ldots,s)\sum_{j=1}^{s}g_{1}(j),\quad s\geq 1.$$

In the case of two-body interaction potential (1.3) operator (4.3) has the form

$$(\mathfrak{B}^{+}g)_{s}(Y)=\mathcal{N}_{s}(Y)g_{s}(Y)+\sum_{j_{1}\neq j_{2}=1}^{s}\mathcal{N}^{(2)}_{\mathrm{int}}(j_{1},j_{2})g_{s-1}(Y\backslash\{j_{1}\}),\quad s\geq 1, \tag{4.6}$$

where the operator $\mathcal{N}_{\mathrm{int}}^{(2)}$ is defined by formula (1.11) for $n = 2$. If $\mathcal{H} = L^2(\mathbb{R}^3)$ in terms of kernels of operators g_s, $s \geq 1$, for expression (4.6) we have

$$(\mathfrak{B}^+ g)_s(q_1, \ldots, q_s; q'_1, \ldots, q'_s) =$$
$$-\frac{i}{\hbar}\Big(-\frac{\hbar^2}{2}\sum_{i=1}^{s}(-\Delta_{q_i} + \Delta_{q'_i}) + \sum_{1=i<j}^{s}\big(\Phi^{(2)}(q'_i - q'_j) - \Phi^{(2)}(q_i - q_j)\big)\Big) g_s(q_1, \ldots, q_s; q'_1, \ldots, q'_s) -$$
$$\frac{i}{\hbar}\sum_{1=i\neq j}^{s}\big(\Phi^{(2)}(q'_i - q'_j) - \Phi^{(2)}(q_i - q_j)\big) g_{s-1}(q_1, \ldots, \overset{j}{\vee}, \ldots, q_s; q'_1, \ldots, \overset{j}{\vee}, \ldots, q'_s),$$

where $(q_1, \ldots, \overset{j}{\vee}, \ldots, q_s) \equiv (q_1, \ldots, q_{j-1}, q_{j+1}, \ldots, q_s)$. This expression for a system of classical particles is defined as a generator of the dual BBGKY hierarchy stated in [1],[25].

In the paper [25] for classical systems of particles an equivalent representation of group (4.2) was used, namely for the case under consideration group the $U^+(t)$ has the following representation

$$U^+(t) = \mathcal{G}(t) + \sum_{n=1}^{\infty}\frac{1}{n!}\big[\ldots\big[\mathcal{G}(t), \underbrace{\mathfrak{a}^+\big], \ldots, \mathfrak{a}^+}_{\text{n-times}}\big] = e^{-\mathfrak{a}^+}\mathcal{G}(t)e^{\mathfrak{a}^+}, \tag{4.7}$$

where $\big[\,.\,,\,.\,\big]$ is a commutator, the operator $\mathfrak{a}^+$ (an analog of the creation operator) is defined on the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ by the formula

$$\big(\mathfrak{a}^+ g\big)_s(Y) := \sum_{j=1}^{s} g_{s-1}(Y\backslash\{j\}), \tag{4.8}$$

the operators $e^{\pm\mathfrak{a}^+}$ are defined on the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ by the expansions

$$\big(e^{\pm\mathfrak{a}^+} g\big)_s(Y) = \sum_{n=0}^{s}\frac{(\pm 1)^n}{n!}\sum_{j_1\neq\ldots\neq j_n=1}^{s} g_{s-n}\big(Y\backslash\{j_1, \ldots, j_n\}\big), \quad s \geq 1.$$

and the group of operators $\mathcal{G}(t)$ is defined by expression (1.6).

Representation (4.7) is true in consequence of definition (4.8) of the operator $\mathfrak{a}^+$ and the validity of the equality

$$\begin{aligned}&\Big(\frac{1}{n!}\big[\ldots\big[\mathcal{G}(t), \underbrace{\mathfrak{a}^+\big], \ldots, \mathfrak{a}^+}_{\text{n-times}}\big] g\Big)_s(Y) = \\ &\frac{1}{(s-n)!}\sum_{j_1\neq\ldots\neq j_{s-n}=1}^{s}\mathfrak{A}^+_{1+n}\big(t, (Y\backslash X)_1, X\big)\, g_{s-n}(Y\backslash X), \quad s \geq 1,\end{aligned} \tag{4.9}$$

where $X \equiv Y\backslash\{j_1, \ldots, j_{s-n}\}$ and $Y\backslash X = (j_1, \ldots, j_{s-n})$. For example, if $n = 1$ we have

$$\big(\big[\mathcal{G}(t), \mathfrak{a}^+\big] g\big)_s(Y) = \sum_{j=1}^{s}\big(\mathcal{G}_s(t, Y) - \mathcal{G}_{s-1}(t, Y\backslash\{j\})\big) g_{s-1}(Y\backslash\{j\}) =$$
$$\sum_{j=1}^{s}\mathfrak{A}^+_2\big(t, (Y\backslash\{j\})_1, j\big)\, g_{s-1}(Y\backslash\{j\})$$

and since identity (4.5) holds we obtain equality (4.9).

We can obtain one more representation of the group $U^+(t)$, if we express the dual cumulants $\mathfrak{A}^+_{1+n}(t),\ n\geq 1$, of groups (1.6) with respect to the $1st$-order and $2nd$-order dual cumulants. In fact it holds

$$\mathfrak{A}^+_{1+n}\big(t,(Y\backslash X)_1,X\big)=\sum_{\substack{Z\subset X,\\ Z\neq\emptyset}}\mathfrak{A}^+_2\big(t,Y\backslash X,Z\big)\sum_{\mathrm{P}:\,X\backslash Z=\bigcup_i X_i}(-1)^{|\mathrm{P}|}\,|\mathrm{P}|!\,\prod\nolimits_{i=1}^{|\mathrm{P}|}\mathfrak{A}^+_1\big(t,X_i\big),$$

where $\sum_{\substack{Z\subset X,\\ Z\neq\emptyset}}$ is a sum over all nonempty subsets $Z\subset X$ of the set X. Then taking into account an identity

$$\sum_{\mathrm{P}:\,X\backslash Z=\bigcup_i X_i}(-1)^{|\mathrm{P}|}\,|\mathrm{P}|!\,\prod\nolimits_{i=1}^{|\mathrm{P}|}\mathfrak{A}^+_1\big(t,X_i\big)g_{s-n}(Y\backslash X)=\sum_{\mathrm{P}:\,X\backslash Z=\bigcup_i X_i}(-1)^{|\mathrm{P}|}\,|\mathrm{P}|!\,g_{s-n}(Y\backslash X)$$

and equality (3.9) we get the following representation of the group $U^+(t)$

$$\begin{aligned}&\big(U^+(t)g\big)_s(Y)=\mathfrak{A}^+_1\big(t,Y\big)g_s(Y)+\\ &\sum_{n=1}^{s}\frac{1}{(s-n)!}\sum_{j_1\neq\ldots\neq j_{s-n}=1}^{s}\sum_{\substack{Z\subset X,\\ Z\neq\emptyset}}(-1)^{|X\backslash Z|}\,\mathfrak{A}^+_2\big(t,Y\backslash X,Z\big)\,g_{s-n}(Y\backslash X).\end{aligned}$$

An infinitesimal generator of group (4.7) has the following form

$$\mathfrak{B}^+=\mathcal{N}+\sum_{n=1}^{\infty}\frac{1}{n!}\big[\ldots\big[\mathcal{N},\underbrace{\mathfrak{a}^+\big],\ldots,\mathfrak{a}^+}_{\text{n-times}}\big]=e^{-\mathfrak{a}^+}\mathcal{N}e^{\mathfrak{a}^+}. \tag{4.10}$$

Representation (4.10) is true in consequence of definition (4.8) of the operator $\mathfrak{a}^+$ and the validity of an identity

$$\big(\big[\ldots\big[\mathcal{N},\underbrace{\mathfrak{a}^+\big],\ldots,\mathfrak{a}^+}_{\text{n-times}}\big]g\big)_s(Y)=\sum_{k=n+1}^{s}\frac{1}{(k-n)!}\sum_{j_1\neq\ldots\neq j_k=1}^{s}\mathcal{N}^{(k)}_{\mathrm{int}}(j_1,\ldots,j_k)g_{s-n}\big(Y\backslash\{j_1,\ldots,j_n\}\big),$$

which in the case of a two-body interaction potential reduces to the following one

$$\big(\big[\mathcal{N},\mathfrak{a}^+\big]g\big)_s(Y)=\sum_{j_1\neq j_2=1}^{s}\mathcal{N}^{(2)}_{\mathrm{int}}(j_1,j_2)g_{s-1}\big(Y\backslash\{j_1\}\big).$$

5 Conclusion

The concept of cumulants (1.9) of groups (1.1) of the von Neumann equations forms the basis of group expansions for quantum evolution equations, namely, the von Neumann hierarchy for correlation operators [23], as well as the BBGKY hierarchy for s-particle density operators [24] and the dual BBGKY hierarchy [26]. In the case of quantum systems of particles obeying Fermi or Bose statistics groups (2.1),(3.2) and (4.2) have different structures. The analysis of these cases will be given in a separate paper.

We have stated the properties of groups (2.1) and (3.2) on the space $\mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$ and dual group (4.2) on $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$. To describe the evolution of infinitely many particles [1] it is necessary to define the one-parametric family of operators (3.2) on more general spaces than $\mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$, for example, on the space of sequences of bounded operators containing the equilibrium states [22]. For dual group (4.2) the problem lies in the definition of functional (1.5) for operators from the corresponding spaces. In both these cases every term of the corresponding expansions contains the divergent traces [1],[24],[25] and the analysis of such a question for quantum systems remains an open problem.

On the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ one-parametric mapping (4.2) is not a strong continuous group. The group $\{U^+(t)\}_{t\in\mathbb{R}}$ of operators (4.2) defined on the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ is dual to the strong continuous group $\{U(t)\}_{t\in\mathbb{R}}$ of operators (3.2) for the BBGKY hierarchy defined on the space $\mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$ and the fact that one is a C^*_0-group follows also from general theorems about properties of the dual semigroups [13],[17].

As mention above the group $\{\mathcal{G}(-t)\}_{t\in\mathbb{R}}$ of operators (1.1) preserves positivity [16],[18]. The same property must be valid for the group $\{U(t)\}_{t\in\mathbb{R}}$ of operators (3.2) for the BBGKY hierarchy, but how to prove this property one is an open problem.

We have constructed infinitesimal generators (2.2), (3.3) on the subspace $\mathfrak{L}^1_{\alpha,0} \subset \mathfrak{L}^1_\alpha(\mathcal{F}_\mathcal{H})$ and generator (4.3) on $\mathcal{D}(\mathfrak{B}^+) \subset \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$. The question of how to define the domains of the definition $\mathcal{D}(\mathfrak{N}), \mathcal{D}(\mathfrak{B})$ and $\mathcal{D}(\mathfrak{B}^+)$ of corresponding generators (2.2), (3.3) and (4.3) is an open problem [13],[18].

References

[1] C.Cercignani, V.I.Gerasimenko and D.Ya.Petrina, *Many-Particle Dynamics and Kinetic Equations.* Kluwer Acad. Publ., 1997.

[2] A.Arnold, *Mathematical properties of quantum evolution equations.* Lect. Notes in Math. **1946**, Springer, 2008.

[3] H.Spohn, *Kinetic equations for quantum many-particle systems.* arXiv:0706.0807v1, 2007.

[4] D.Benedetto, F.Castella, R.Esposito and M.Pulvirenti, *A short review on the derivation of the nonlinear quantum Boltzmann equations.* Commun. Math. Sci. **5**, (2007), 55–71.

[5] R.Adami, F.Golse and A.Teta, *Rigorous Derivation of the Cubic NLS in Dimension One.* J. Stat. Phys. **127**, (6), (2007), 1193-1220.

[6] L.Erdős, M.Salmhofer and H.-T.Yau, *On quantum Boltzmann equation.* J. Stat. Phys. **116**, (116), (2004), 367-380.

[7] L.Erdős, B.Schlein and H.-T.Yau, *Derivation of the cubic non-linear Schröodinger equation from quantum dynamics of many-body systems.* Invent. Math. **167**, (3), (2007), 515-614.

[8] L.Erdős and B.Schlein, *Quantum dynamics with mean field interactions: a new approach.* arXiv:0804.3774v1, 2008.

[9] J.Fröhlich, S.Graffi and S.Schwarz, *Mean-field- and classical limit of many-body Schrödinger dynamics for bosons.* Commun. Math. Phys., **271**, (2007), 681-697.

[10] R.Dautray and J.L.Lions, *Mathematical Analysis and Numerical Methods for Science and Technology.* **5**, Springer-Verlag, 1992.

[11] D.Ya.Petrina, *Mathematical Foundations of Quantum Statistical Mechanics. Continuous Systems.* Kluwer, 1995.

[12] F.A.Berezin and M.A.Shoubin, *Schrödinger Equation.* Kluwer, 1991.

[13] O.Bratelli and D.W.Robinson, *Operator Algebras and Quantum Statistical Mechanics.* **1**, Springer-Verlag, 1979.

[14] T.Kato, *Perturbation Theory for Linear Operators.* Springer-Verlag, 1995.

[15] A.Bellini-Morante and A.C.McBride, *Applied Nonlinear Semigroups.* John Wiley and Sons, 1998.

[16] J.Banasiak and L.Arlotti, *Perturbations of Positive Semigroups with Applications.* Springer, 2006.

[17] A.Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations.* Springer-Verlag, 1983.

[18] R.Aliki and K.Lendi, *Quantum dynamical semigroups and applications.* Lect. Notes in Phys. **286**, Springer, 1987.

[19] P.A.Markowich, *On the equivalence of the Schrödinger and the quantum Liouville equations.* Math. Meth. Appl. Sci. **11**, (1989), 459-469.

[20] V.O.Shtyk, *On the solutions of the nonlinear Liouville hierarchy.* J. Phys. A: Math. Theor. **40**, (2007), 9733-9742.

[21] M.S.Green, *Boltzmann equation from the statistical mechanical point of view.* J. Chem. Phys. **25**, (5), (1956), 836-855.

[22] J.Ginibre, *Some applications of functional integrations in statistical mechanics.* In *Statistical Mechanics and Quantum Field Theory.* (Eds. S.De Witt and R.Stora, Gordon and Breach), (1971), 329-427.

[23] V.I.Gerasimenko and V.O.Shtyk, *Evolution of correlations of quantum many-particle systems.* J. Stat. Mech. (3), (2008), P03007, 24p.

[24] V.I.Gerasimenko and V.O.Shtyk, *Initial-value problem for the Bogolyubov hierarchy for quantum systems of particles.* Ukrain. Math. J. **58**, (9), (2006), 1329-1346.

[25] G.Borgioli and V.Gerasimenko, *The dual BBGKY hierarchy for the evolution of observables.* Riv. Mat. Univ. Parma. **4**, (2001), 251-267.

[26] V.I.Gerasimenko and T.V.Ryabukha, *Cumulant representation of solutions of the BBGKY hierarchy of equations.* Ukrain. Math. J. **54**, (10), (2002), 1583-1601.

[27] V.I.Gerasimenko, T.V.Ryabukha and M.O.Stashenko, *On the structure of expansions for the BBGKY hierarchy solutions.* J. Phys. A: Math. Gen. **37**, (2004), 9861-9872.